



Serum Uric Acid Level, Alzheimer-type Dementia and Variation

Serum Ürik Asit Düzeyi, Alzheimer Tipi Demans ve Varyasyon

Beuy Joob¹, Viroj Wiwanitkit²

¹Sanitation Medical Academic Center, Bangkok, Thailand

²Dr. DY Patil University, Pune, India

Keywords: Uric acid, Alzheimer, dementia, variation

Anahtar Kelimeler: Ürik asit, Alzheimer, demans, varyasyon

Dear Editor,

We read the publication entitled “Assessment of Serum Uric Acid (UA) Levels According to Sex and Stage for Patients with Alzheimer-type Dementia” with great interest (1). Yazar and Yazar (1) concluded that “Our study supports the hypothesis that the decrease detected in serum UA concentrations in parallel with the progression of disease stages is directly related to the pathogenesis of AD”. We agree that there might be some possible pathophysiologic mechanisms that might affect serum UA levels in AD. Nevertheless, there are many considerations regarding the variation of UA based on laboratory medicine principles. First, Yazar and Yazar (1) had the UA measurement in two different laboratories from two hospitals. A variation of UA determination can be seen. The differences of measured UA levels due to different laboratory technicians as well as different laboratories are well documented in the literature (2,3). In the present report, the inter-personal and inter-laboratory variations can be expected. In addition, there is also a seasonal variation of UA levels (4). These factors might affect the observation in the report by Yazar and Yazar (1).

Authorship Contributions

Surgical and Medical Practices: BJ., VW., Concept: BJ., VW., Design: BJ., VW., Data Collection or Processing: BJ., VW.,

Analysis or Interpretation: BJ., VW., Literature Search: BJ., VW., Writing: BJ., VW.

Ethics

Informed Consent: Not applicable.

Peer-review: Externally peer-reviewed.

Financial Disclosure: The authors declared that this study received no financial support.

References

1. Yazar T, Yazar HO. Assessment of serum uric acid levels according to sex and stage for patients with Alzheimer-type dementia. Turk J Neurol 2019;25:76-81.
2. Jiang Y, Liu O, Xu G. Assessment of the trueness and inter-laboratory precision of routine uric acid assays using 4 frozen pooled serum samples measured by the Japan Society of Clinical Chemistry's HPLC method. Ann Lab Med 2014;34:104-110.
3. Chen MS, Liao CM, Wu MH, Lin CM. Improvement of the quality of work in a biochemistry laboratory via measurement system analysis. Chin J Physiol 2016;59:293-299.
4. Murciano Revert J, Martínez-Lahuerta JJ, Aleixandre Porcar L. Seasonal change in blood concentration of uric acid and its potential clinical implications. Aten Primaria 2000;26:468-471.

Address for Correspondence/Yazışma Adresi: Beuy Joob MD, Sanitation Medical Academic Center, Bangkok, Thailand

Phone: +6624658292 E-mail: beuyjoob@hotmail.com ORCID: orcid.org/0000-0002-5281-0369

Received/Geliş Tarihi: 20.06.2019 **Accepted/Kabul Tarihi:** 11.11.2019

©Copyright 2020 by Turkish Neurological Society
Turkish Journal of Neurology published by Galenos Publishing House.